



Contribution ID: 10

Type: not specified

Online diagnostics for high-intensity laser-plasma radiation characterization

Tuesday, 29 October 2024 10:10 (25 minutes)

High-intensity laser-plasma interactions are pivotal in advanced scientific research, with applications spanning fundamental physics to medicine. The ELIMAIA (ELI Multidisciplinary Applications of laser-Ion Acceleration) [1-3] beamline at the ELI Beamlines facility, part of the Extreme Light Infrastructure ERIC, offers a cutting-edge platform for studying laser-driven ion acceleration at high repetition rates. For such a beamline, real-time measurements and analysis of the laser-produced radiation are essential characterizing the laser-plasma interaction.

We describe a comprehensive set of online diagnostics developed at the ELIMAIA beamline, enabling immediate and precise analysis of radiation and secondary sources generated from high-intensity laser-plasma interactions. The key diagnostic tools include ion detectors such as online Thomson parabola and semiconductor detectors operating in the Time-of-Flight regime, laser beam diagnostics, and X-ray spectrometers. These tools are integrated with a real-time data acquisition and analysis system.

The diagnostic set was successfully tested in multiple internal and user experiments at ELIMAIA including high repetition rate tests. It has proven invaluable for optimizing ion acceleration and monitoring the quality of laser-target interactions.

Our results highlight the importance of integrating advanced diagnostics in high-intensity laser-plasma research, driving forward the capabilities of next-generation laser facilities.

[1] D. Margarone, “ELIMAIA: A laser-driven ion accelerator for multidisciplinary applications.” *Quantum Beam Science* 2.2 (2018): 8.

[2] F. Schillaci, “The ELIMAIA laser-plasma ion accelerator: Technological commissioning and perspectives.” *Quantum Beam Science* 6.4 (2022): 30.

[3] L. Giuffrida, et al “The ELIMAIA laser-plasma Ion Accelerator: a user platform of high-repetition-rate, ultrahigh intensity, and high stability”, in preparation.

Primary authors: Dr ISTOKSKAIA, Valeria (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr PSIKAL, Jan (ELI Beamlines facility, Extreme Infrastructure ERIC; Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague); Dr SCHILLACI, Francesco (ELI Beamlines facility, Extreme Infrastructure ERIC); Mr STANCEK, Stanislav (ELI Beamlines facility, Extreme Infrastructure ERIC; Joint Laboratory of Optics of Palacky University, Institute of Physics of Academy of Sciences of the Czech Republic, Faculty of Science, Palacky University); Mr TOSCA, Marco (ELI Beamlines facility, Extreme Infrastructure ERIC; Charles University, Faculty of Mathematics and Physics, Department of Macromolecular Physics); Dr TRYUS, Maksym (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr VELYHAN, Andriy (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr MARGARONE, Daniele (ELI Beamlines facility, Extreme Infrastructure ERIC; Centre for Light-Matter Interactions, School of Mathematics and Physics, Queen’s University Belfast); Dr GIUFFRIDA, Lorenzo (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr CHAGOVETS, Timofej (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr GAMAIUNOVA, Nina (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr GREPL, Filip (ELI Beamlines facility, Extreme Infrastructure ERIC); Dr GREPLOVA ZAKOVA, Martina (ELI Beamlines facility, Extreme Infrastructure ERIC); Mr HADJIKYRIACOU, Arsenios (ELI Beamlines facility, Extreme Infrastructure ERIC; Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague); Dr KANTARELOU, Vasiliki (ELI Beamlines facility, Extreme Infrastructure ERIC)

Presenter: Dr ISTOKSKAIA, Valeria (ELI Beamlines facility, Extreme Infrastructure ERIC)

Session Classification: Oral contributions I